

Combining Biofeedback with Stress Management Interventions: A Systematic Review of Physiological and Psychological Effects

Nele A. J. De Witte, Inez Buyck & Tom Van Daele

Applied Psychophysiology and Biofeedback

In association with the Association for Applied Psychophysiology and Biofeedback

ISSN 1090-0586

Appl Psychophysiol Biofeedback
DOI 10.1007/s10484-018-09427-7



Volume 38, Number 3

Applied Psychophysiology and Biofeedback

 Springer

Available
online
www.springerlink.com

applied
psychophysiology
& biofeedback
aapb
an international society
for research, research
with, and education

 Springer

Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media, LLC, part of Springer Nature. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".



Combining Biofeedback with Stress Management Interventions: A Systematic Review of Physiological and Psychological Effects

Nele A. J. De Witte¹ · Inez Buyck^{1,2} · Tom Van Daele¹

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

Current mental healthcare systems experience difficulties meeting the challenges of a growing population with elevated stress symptoms. Outpatient stress management interventions have already proven to be effective in routine care and recent technological advances now allow to expand such interventions, for example by adding a physiological component like biofeedback. Adding biofeedback to stress management interventions appears promising, but there is a lack of insight into the general conceptualization and evaluation of the resulting interventions, both in relation to psychological and physiological stress indicators. A comprehensive literature search was performed to investigate stress management interventions with a biofeedback component. This systematic review provides an overview of these interventions and explores to what extent they can improve both physiological and psychological indicators of stress. Fourteen RCTs were included. A large diversity was observed in intervention design and effectiveness. Nevertheless, there is preliminary evidence that the use of biofeedback can improve both physiological and psychological indicators of stress. Biofeedback could provide an accessible and low-cost addition to stress interventions. Further research into the effectiveness of different components of biofeedback interventions is needed.

Keywords Biofeedback · Stress reduction · mHealth · Effectiveness · Heart rate variability

Introduction

Stress-related problems are becoming increasingly common and are associated with large health risks (Van Daele et al. 2012). There is a need for interventions that can capture and improve the complex interplay between physiological and psychological stress processes in daily life. Technological advances allow existing psychophysiological laboratory designs for stress management to be modified for use outside of the lab, where mHealth applications could provide immediate feedback in a way that has not yet been possible in traditional mental healthcare (Williams 2016). mHealth refers to the use of mobile information and communication technology, such as mobile computers, medical sensors,

and wearable devices in healthcare (Istepanian et al. 2004). However, prior to making the translation from experimental to clinical applications, research into the underlying mechanisms providing effectiveness, both on a psychological and a physiological level, is needed to establish which interventions have the largest potential for broad application.

Stress is a complex phenomenon that is triggered by a psychological or physical threat to homeostasis and consists of a variety of psychological, behavioral, and physiological responses (Bali and Jaggi 2015). There are many different potential sources of stress, such as work-related stress, stress due to family conditions or stress associated with medical or physical illness (such as chronic pain). Experiencing mild stress is not maladaptive or unhealthy in itself, but experiencing high levels of stress without sufficient recovery is a substantial health risk. Elevated and prolonged experience of stress is a common problem, which has a negative impact on both mental and physical health, and is associated with large economic costs (American Psychiatric Association 2016; Cooper and Dewe 2008). Current mental healthcare systems have difficulties meeting the challenges of a growing population of individuals with elevated stress. A recent

✉ Nele A. J. De Witte
nele.dw@thomasmore.be

¹ Expertise Unit Psychology, Technology & Society, Thomas More University of Applied Sciences, Molenstraat 8, 2018 Antwerp, Belgium

² LUCAS, Centre for Care Research and Consultancy, KU Leuven, Leuven, Belgium

survey showed that 75% of Americans had experienced at least one symptom of stress (e.g., feeling nervous, being irritable or experiencing fatigue) in the past month (American Psychological Association 2017). This is a 4% increase as compared to the previous measurement in 2016.

Stressful situations can evoke changes in heart rate, heart rate variability (HRV), blood pressure, electrodermal activity (EDA), and breathing rate (Chrousos and Gold 1992; Jarczok et al. 2013; Lin et al. 2011). Such physiological parameters can be a useful addition to self-report data, since these do not suffer from reporting or social desirability biases. An example of an intervention strategy strongly based on physiological processes is biofeedback. Biofeedback works to help clients improve their performance and health by gaining voluntary control over real-time physiological processes, such as HRV or EDA (Dillon et al. 2016). Formerly, a large disadvantage of interventions based on physiology was the need for large and expensive devices, which reduced mobility and therefore generalizability outside of the lab. However, the advance of mHealth can reduce the cost and increase the impact of psychophysiological interventions. Wearables have the potential to non-invasively collect behavioral and physiological data and thereby provide additional information that can guide treatment.

The review of Schoenberg and David (2014) shows that biofeedback can be a successful intervention to treat psychiatric symptoms. Biofeedback appears especially relevant when maladaptive physiological mechanisms (such as heightened autonomic nervous system activity) are at play, which is the case in stress experiences. The mechanisms of change in psychophysiological interventions are still largely unknown. Wheat and Larkin (2010) show that HRV biofeedback can induce significant changes in physiological parameters in different medical and psychiatric disorders. However, very little research directly assesses the relationship between physiological and psychological outcomes of biofeedback. It remains unclear whether physiological changes are a prerequisite to psychological effects of such psychophysiological interventions.

As highlighted in the previous paragraphs, psychophysiological parameters could be relevant indicators for mental health, with the required technology to measure these parameters becoming increasingly low-cost and less invasive. As such technology now allows to (pro)actively monitor or intervene outside of highly controlled lab settings, the question arises to what extent there is already evidence available on the use and added value of psychophysiological interventions in mental healthcare. The current systematic review focuses on the question whether there is any evidence for using biofeedback in the context of stress management interventions. Additionally, the association between physiological and psychological outcomes of these interventions is explored.

Method

A search of the literature was performed using broad keywords such as “*physiolog**” and “*psychophysiolog**” combined with “*intervention*” and “*stress*”. This resulted in a striking amount of articles, but it became evident that controlled research into psychophysiological interventions for stress was mostly confined to biofeedback. A closer inspection of these manuscripts also revealed that a substantial subgroup of results focused on stress in the context of posttraumatic stress disorder (PTSD) or urinary stress incontinence. The current review is focused on the management of ongoing psychological stressors (e.g., demanding occupations or chronic pain) and, therefore, PTSD and urinary stress incontinence are out of scope. Consequently, the focus of the literature search was narrowed by use of the PICO-TS model to improve the relevance of the results and maximize comparability between studies. The current review aims to investigate the general population (P) with potential stress complaints. The intervention (I) is the main focus of the review and consists of biofeedback, either stand-alone or in addition to other stress management interventions. There was a special interest in mobile health interventions (using wearable devices) when they were available. There were no requirements pertaining to the control condition (C), but the reported outcomes (O) did need to consist of both physiological and psychological indices. No timings (T) were specified and with regard to study design (S), only randomized controlled trials were included.

A literature search of studies published until September 2017 was set up through search engines PubMed (MEDLINE), Web of Science, PsycINFO, and Cochrane Library database. The systematic search implemented the following string of keywords: (biofeedback OR wearable) AND stress NOT (urinary or posttraumatic or PTSD) AND (interven* OR treat* OR ther* OR manag*). Articles were selected for review based on three inclusion criteria: (1) studies implemented psychophysiological indices in a stress management intervention; (2) study designs were randomized controlled trials with a valid psychological as well as physiological measure of stress before and after the intervention; (3) articles needed to be published in an international, peer-reviewed journal and written in English. Studies were excluded when psychophysiological indices were used for diagnostic purposes only.

In a first step, two authors (IB and TVD) independently evaluated titles and abstracts for eligibility criteria. After resolving disagreements through discussion, the same authors were involved in full text evaluation. Subsequently, reference sections of the included studies were manually inspected for additional articles, yet no supplemental

studies could be included. The studies that were found showed large diversity in the design and outcome variables, thereby eliminating the possibility of performing a meta-analysis. Relevant data on the samples, interventions and outcomes variables was extracted from the articles and summarized in tables. Additionally, a Fisher's Exact test was performed to investigate the association between the occurrence of physiological and psychological outcomes.

A risk of bias analysis was conducted using the Cochrane Risk of Bias Tool (Higgins et al. 2011). The included studies were screened for selection bias, performance bias, detection bias, attrition bias, and reporting bias. Studies were considered to be highly susceptible for selection bias if violations were detected in the random assignment of participants to intervention or control groups. Selection bias was regarded low if adequate procedures regarding random group assignment were conducted, resulting in comparable groups to be investigated. Performance bias was considered to be high when there was no blinding (or masking) of study participants and personnel. Although difficult to set up in biofeedback studies, such a procedure reduces the risk that mere knowledge of the offered intervention affects outcome. When successful blinding was undertaken, this bias was considered low. Detection bias was considered to be high when researchers analyzing the data were aware of the intervention allocated to participants. Contrary, if the


blinding of these outcome assessors was assured, risk of bias was considered to be low. A high risk for attrition bias was considered to be present in case of incomplete outcome data from the analyses, usually due to drop-out of participants. Studies were regarded to be low in risk for attrition bias if the data of each outcome measure were complete. Selective outcome reporting, i.e., not reporting the results of one of the outcome measures, was considered an indication of reporting bias. Studies were evaluated low in reporting bias if the results of all outcome measures were reported, in comparison to published study protocols. In all risk of bias analyses, insufficient details on the respective aspect of the study methodology resulted in a statement of unclear risk of bias.

Results

Study Selection

The search generated 2056 potentially relevant articles (Fig. 1), which were reduced to 1718 articles after removal of duplications. A total of 1605 articles were eliminated following title and abstract screening. Finally, after reading the full text another 85 articles were excluded because they did

Fig. 1 Flow chart of the search strategy



not fulfil inclusion criteria. In all, this resulted in the inclusion of 14 articles for review.

Risk of Bias Evaluation

All studies randomized participants across study conditions, resulting in satisfactory conditions in terms of selection bias (Table 1). Performance bias was often considered to be present, as clear-cut control conditions were frequently used, with only a single study opting for a double blind design. Furthermore, it was difficult to assess whether any detection bias occurred, as most studies failed to provide input on whether or not outcome assessors were blinded. Attrition bias did seem to be present in several studies. This is typically due to drop-out of participants during the intervention or assessment, resulting in incomplete compilation of data. Drop-out rate itself was, however, well documented in most studies. Finally, no study protocols appeared to have been pre-published, which is however, only a fairly recent practice. Most articles did make use of a non-published protocol, or provided a comprehensive overview of their study results. It was therefore concluded that overall there seemed to be little evidence of reporting bias.

Study Focus and Sample

As Table 2 illustrates, the majority of included studies focused specifically on stress management or stress reduction, although one study was aimed at general self-control (Mackay et al. 2015). The stress interventions were implemented in a healthy population (Dillon et al. 2016; Kotozaki et al. 2014; Murphy 1984; Whited et al. 2014), a population with demanding job characteristics (Allen and Blanchard

1980; Lemaire et al. 2011; McCraty et al. 2009; Sutarto et al. 2012), or in a medical context, targeting patients with chronic pain (Berry et al. 2014; Hallman et al. 2011), multiple sclerosis (Mackay et al. 2015), heart disease (Nolan et al. 2005), obesity (Teufel et al. 2013), and pregnant women at risk for preterm labor (Siepmann et al. 2014). In total, the included studies comprised 488 participants, of which 51% was female ($n=250$). Participants were between 18 and 60 years old (pooled $M=39.72$, pooled $SD=7.61$; Allen and Blanchard (1980) were not included in the pooled statistics since they only reported age range).

Biofeedback Intervention

Nine studies implemented stress interventions that relied on variations of biofeedback on heart rate (Kotozaki et al. 2014) and heart rate coherence or HRV (Berry et al. 2014; Hallman et al. 2011; Lemaire et al. 2011; McCraty et al. 2009; Nolan et al. 2005; Siepmann et al. 2014; Sutarto et al. 2012; Whited et al. 2014; Table 3). Two recent studies used EDA as the leading physiological parameter for biofeedback (Dillon et al. 2016; Teufel et al. 2013), while in two older studies, feedback was provided on electromyographic (EMG) activity (Allen and Blanchard 1980; Murphy 1984). One final study applied breathing rate and muscle tension biofeedback (Mackay et al. 2015).

The duration of the biofeedback interventions ranged from one single session (Dillon et al. 2016) to an intervention that was spread out over the course of 12 weeks (McCraty et al. 2009), with most interventions being implemented for approximately 4 weeks (Berry et al. 2014; Kotozaki et al. 2014; Lemaire et al. 2011; Murphy 1984; Nolan et al. 2005). The length and number of sessions showed large

Table 1 Overview of risk of bias, ranging from low (–), over unclear (0) to high (+) risk of bias

Study	Selection bias	Performance bias	Detection bias	Attrition bias	Reporting bias
1. Allen and Blanchard (1980)	–	+	–	–	–
2. Berry et al. (2014)	–	+	0	–	–
3. Dillon et al. (2016)	–	0	0	–	–
4. Hallman et al. (2011)	–	0	–	+	–
5. Kotozaki et al. (2014)	–	–	0	–	+
6. Lemaire et al. (2011)	–	+	+	+	–
7. Mackay et al. (2015)	–	0	0	–	–
8. McCraty et al. (2009)	–	+	0	+	–
9. Murphy (1984)	–	+	0	+	+
10. Nolan et al. (2005)	–	+	0	+	–
11. Siepmann et al. (2014)	–	+	0	+	–
12. Sutarto et al. (2012)	–	+	+	+	–
13. Teufel et al. (2013)	–	+	0	+	–
14. Whited et al. (2014)	–	+	0	+	–

Table 2 Study design features

Study	Study focus	Sample size (# female)	Sample characteristics and mean age (SD)	Physiological measurements	Psychological measurements	Follow-up
1. Allen and Blanchard (1980)	Stress management	30 (8)	Middle-level managers Age range: 40–60	Frontal electro-myography, finger temperature	SSS	6 weeks (SSS)
2. Berry et al. (2014)	Pain and stress management	14 (1)	Veterans with chronic pain 44.63 (6.96)	HRV coherence	PSS	–
3. Dillon et al. (2016)	Stress reduction	50 (32)	Healthy adults 26.7 (5.1)	Heart rate	Visual Analogue Scale for perceived stress	–
4. Hallman et al. (2011)	Autonomic regulation and perceived health, pain, stress and disability	24 (22)	Adults with stress-related chronic neck pain 41.43 (7.07)	HRV during rest, during hand grip test, during cold pressor test, and during deep breathing test	Stress Medicine Symptom Scale	–
5. Kotozaki et al. (2014)	Stress management	30 (0)	Healthy working adults 42.30 (7.90)	Salivary cortisol and voxel-based morphometry	Brief Job Stress Questionnaire	–
6. Lemaire et al. (2011)	Stress management	40 (17)	Physicians 46.3 (8.35)	Heart rate, blood pressure, salivary cortisol	Scale based on PSS and POQA—revised	28 days
7. Mackay et al. (2015)	Self-control	40 (32)	Patients with multiple sclerosis 45.9 (12.42)	Muscle tension, breathing rate	DASS	3 months (DASS)
8. McCraty et al. (2009)	Stress and health risk reduction	75 (23)	Correctional officers 40.00 (7.88)	HRV, blood pressure, blood samples, and saliva samples	POQA	–
9. Murphy (1984)	Stress management	38 (4)	Highway maintenance workers 42 (SD not given)	Forehead electro-myography	Job Stress Scale	3 months
10. Nolan et al. (2005)	Stress and depression reduction	46 (6)	Patients with coronary heart disease 54.52 (1.26)	HRV at baseline, during physical, emotional, and personal stressor, and during recovery	PSS	–
11. Siepmann et al. (2014)	Reduction of stress and preterm birth	48 (48)	Pregnant women prone to preterm labor 29 (5)	Heart rate, HRV	Trier Inventory for the Assessment of Chronic Stress	4 weeks
12. Sutarto et al. (2012)	Stress reduction	36 (36)	Manufacturing operators 36.30 (10.14)	HRV during rest, stressor, and recovery	DASS	–
13. Teufel et al. (2013)	Self-efficacy and stress reduction	30 (30)	Obese adults 48.67 (9.94)	Electrodermal activity to food stimuli	Perceived Stress Questionnaire	3 months
14. Whited et al. (2014)	Stress reduction	27 (23)	Students 22.54 (3.82)	HRV at rest, during stressor and during recovery	PSS	–

Pooled means were calculated for age (in years) when no total group means were provided. Allen and Blanchard (1980) and Murphy (1984) did not report the mean and/or standard deviation of age

DASS Depression Anxiety Stress Scales, HRV heart rate variability, POQA Personal and Organizational Quality Assessment, PSS Perceived Stress Scale, SSS Subjective Stress Scale

Table 3 Overview of the intervention characteristics and results of the included studies

Study	Intervention duration	Experimental intervention	Control condition(s)	Physiological effects	Psychological effects	Conclusion
1. Allen and Blanchard (1980)	6 weeks 55 min/week + Daily practice	EMG BF, progressive relaxation, abdominal breathing, and stress inoculation training	(1) WLC (2) 6 weekly individual and group sessions on stress	Decrease in basal frontal EMG and frontal EMG during stress recovery: INT and C	No effects	Physiological effects but no superior effect of INT
2. Berry et al. (2014)	4 weeks 1 session/week	HRV coherence BF and TAU	TAU	POST HRV coherence: INT > C	POST perceived stress: INT < C	Larger physiological and psychological effects in INT than C
3. Dillon et al. (2016)	1 session of 30 min	Two EDA BF games	Non-biofeedback game	POST HR: INT < C	POST perceived stress: INT < C	Larger physiological and psychological effects in INT than C
4. Hallman et al. (2011)	10 weeks 1 session/week + 15 min daily practice	Resonance HRV BF	Breathing protocol in week 1 and 10, no intermittent treatment	POST HRV during rest, reactivity to stress and recovery from stress: INT > C	No effects	Larger physiological effects in INT than C (but not in all HRV indices)
5. Kotozaki et al. (2014)	4 weeks 5 min/day	HR and CBF BF	No intervention	Decrease in salivary cortisol level: INT > C Increase in grey matter volume of right lateral orbitofrontal cortex and left subgenual anterior cingulate cortex: INT > C	Decrease in tension and stressors of working environment score: INT > C	Larger physiological and psychological effects in INT than C
6. Lemaire et al. (2011)	4 weeks 3 sessions of 5 min/day	HRV coherence BF, rhythmic breathing, self-generated positive emotion, and components control condition	Information brochure and biweekly contact with research assistant (+BF training after 4 weeks)	No effects	Decrease in perceived stress: INT > C Effects INT maintained at FU	Larger psychological effects in INT than C
7. Mackay et al. (2015)	3 weeks 60 min/week + Daily practice	Breathing rate and muscle tension BF and components control condition	Relaxation, mindfulness, social support, and education	No effects	Decrease in stress: INT	Psychological effects but no significant group differences
8. McCraty et al. (2009)	2-day training program + 3 months of daily practice	HRV coherence BF and "Power to Change Performance" program	WLC	Decrease in blood pressure, HR, and dehydroepiandro-sterone (DHEA) and increase in HRV: INT Decrease in cholesterol and glucose: INT and C	No effects	Physiological effects in INT but no significant group differences

Table 3 (continued)

Study	Intervention duration	Experimental intervention	Control condition(s)	Physiological effects	Psychological effects	Conclusion
9. Murphy (1984)	2 weeks 60 min/day + Practice between sessions	Forehead EMG BF	(1) WLC (2) MR	Decrease in EMG: INT > WLC Percentages of EMG reduction at FU: INT > MR > WLC	No effects	Larger physiological effects in INT than C
10. Nolan et al. (2005)	4 weeks 5 sessions of 90 min	Paced breathing guided by HRV BF and cognitive-behavioral training	TAU	POST HRV increase during stress recovery: INT > C	Decrease in stress: INT and C	Larger physiological effects in INT than C, no group difference in psychological effects, inverse association between stress and HRV modulation limited to INT
11. Siepman et al. (2014)	2 weeks 3 sessions of minimum 25 min/week	Paced breathing guided by HRV BF	Passive viewing condition	No effects	FU decrease in chronic stress: INT	Psychological effects in INT at FU but no significant group differences
12. Sutar et al. (2012)	5 weeks 30–50 min/week + 20 min daily practice without device	Paced breathing guided by HRV BF	5 sessions of 20 min physiological monitoring, without (BF) instructions	POST increase in HRV during baseline, stress and recovery: INT > C Decrease in breaths per minute during baseline, stress, and recovery: INT	Decrease in stress: INT (no direct group comparison performed)	Physiological and psychological effects in INT but group differences were not always assessed or significant
13. Teufel et al. (2013)	8 weeks 21 min/week + Daily practice without device	EDA BF (1) specific INT: food photos (2) relaxation INT: landscapes	WLC	POST and FU reduction in EDA; medium effect in specific INT, small effect in relaxation INT	POST reduction in perceived stress: small effect in specific INT FU reduction in perceived stress: large effect in specific INT, small effect in relaxation INT	Physiological and psychological effects in INT but no significant group differences
14. Whited et al. (2014)	4 to 8 weeks 32 min/week + Daily practice	HRV coherence BF with exercises for attention, breathing and inducing positive emotions	No intervention	POST HRV during stress INT > C	No effects	Larger physiological effect in INT than C

BF biofeedback, C control condition, CBF cerebral blood flow, EDA electrodermal activity, EMG electromyography, FU follow-up, HRV heart rate, HRV heart rate variability, INT intervention, MR muscle relaxation, TAU treatment as usual, WLC waiting list control

variability as well. Most studies encouraged daily practice of the learned skills (Allen and Blanchard 1980; Hallman et al. 2011; Kotozaki et al. 2014; Lemaire et al. 2011; Mackay et al. 2015; McCraty et al. 2009; Murphy 1984; Sutarto et al. 2012; Teufel et al. 2013; Whited et al. 2014).

The studies offered biofeedback as a stand-alone intervention (Dillon et al. 2016; Hallman et al. 2011; Kotozaki et al. 2014; Murphy 1984; Siepmann et al. 2014; Sutarto et al. 2012; Teufel et al. 2013) or as part of a broader intervention program for stress management (Allen and Blanchard 1980; Berry et al. 2014; Lemaire et al. 2011; Mackay et al. 2015; McCraty et al. 2009; Nolan et al. 2005; Whited et al. 2014). The training was offered in a controlled laboratory setting (Berry et al. 2014; Dillon et al. 2016; Hallman et al. 2011; Mackay et al. 2015; Nolan et al. 2005; Teufel et al. 2013; Whited et al. 2014), at work (Allen and Blanchard 1980; Murphy 1984; Sutarto et al. 2012), at home (Kotozaki et al. 2014), or outside of the lab without further specifications (Lemaire et al. 2011; McCraty et al. 2009). Siepmann et al. (2014) did not specify the location of the intervention.

Control Condition

The implemented control conditions were also characterized by high levels of diversity. Six studies included passive control groups, specifically a waiting list control group (Allen and Blanchard 1980; McCraty et al. 2009; Murphy 1984; Teufel et al. 2013) or no intervention (Kotozaki et al. 2014; Whited et al. 2014). However, Allen and Blanchard (1980) and Murphy (1984) additionally included a second active control condition. Active control conditions consisted of standard care for stress management (Berry et al. 2014; Nolan et al. 2005), individual and group discussions on stress (Allen and Blanchard 1980), a breathing protocol (Hallman et al. 2011), educational information (Lemaire et al. 2011), muscle relaxation (Murphy 1984), or a combination of relaxation, mindfulness, social support and education (Mackay et al. 2015). Finally, three studies designed a control condition with corresponding features to the biofeedback intervention, such as playing a game (Dillon et al. 2016), presenting the visuals of the experimental intervention at the same frequency and duration, but without instruction (Siepmann et al. 2014), or physiological monitoring without providing biofeedback (Sutarto et al. 2012).

Outcome Measures

Most studies evaluated physiological treatment effects through the measure that was implemented in the biofeedback condition, specifically frontal EMG (Allen and Blanchard 1980; Murphy 1984), EDA (Teufel et al. 2013), muscle tension and breathing rate (Mackay et al. 2015), and HRV (coherence) in a resting state, in response to stress, or

in a relaxation condition (Berry et al. 2014; Hallman et al. 2011; Nolan et al. 2005; Siepmann et al. 2014; Sutarto et al. 2012; Whited et al. 2014). However, four studies also used different physiological modalities in the outcome measures as compared to the biofeedback intervention, such as Dillon et al. (2016) who assessed the effect of EDA biofeedback on heart rate, Kotozaki et al. (2014) who used salivary cortisol and grey matter volumes as outcome measures for heart rate and cerebral blood flow biofeedback, and Lemaire et al. (2011) and McCraty et al. (2009) who implemented HRV coherence biofeedback and measured heart rate, blood pressure and several molecules (such as cortisol). The included studies measured HRV in both the frequency and time domain, with most studies implementing multiple HRV indices. With regard to the measurement of psychological treatment effects, there is not a golden standard for the measurement of stress. The Perceived Stress Scale (PSS; Cohen et al. 1983) was used three times (Berry et al. 2014; Nolan et al. 2005; Whited et al. 2014) and the Depression Anxiety Stress Scales (DASS; Lovibond and Lovibond 1995) was included twice (Mackay et al. 2015; Sutarto et al. 2012), all of the other studies implemented different assessments.

Effects of the Intervention

Berry et al. (2014), Dillon et al. (2016), and Kotozaki et al. (2014) observed beneficial effects of biofeedback for stress management as compared to the control condition in both the physiological and psychological domain. Five additional studies observed a significant physiological effect of the intervention, as compared to the control condition (Hallman et al. 2011; Murphy 1984; Nolan et al. 2005; Sutarto et al. 2012; Whited et al. 2014). Finally, one study observed improved psychological outcomes in the biofeedback intervention, as compared the control condition (Lemaire et al. 2011). HRV was the most commonly affected physiological outcome variable, specifically HRV in recovery from stress (Hallman et al. 2011; Nolan et al. 2005; Sutarto et al. 2012; Whited et al. 2014) or at rest (Hallman et al. 2011).

A follow-up measurement was included in five studies. Teufel et al. (2013) observed that both psychological and physiological effects remained or even improved until 3 months follow-up. Lemaire et al. (2011) observed maintained psychological effects at 28 days follow-up and Murphy (1984) detected maintained physiological effects at three months follow-up. Siepmann et al. (2014) did not measure stress immediately after the intervention but did find a reduction in stress at 4 weeks follow-up. Finally, Allen and Blanchard (1980) did not find an effect of biofeedback on self-reported stress immediately after the intervention or at 6 weeks follow-up (these authors did not include a physiological follow-up measurement).

Association Between Physiological and Psychological Effects

Table 4 provides an overview of the occurrence of significant psychological and physiological outcomes, as compared to the control condition, in the fourteen included studies. The results showed that 38% of studies observing beneficial physiological outcomes also observed reduced stress levels and 75% of studies with reduced stress levels also observed improved physiological outcomes, a difference that was not significant ($p = .580$, two tailed Fisher's Exact test).

Effectiveness Related to Different Components of the Biofeedback Intervention and Sample

The included studies showed large diversity in the characteristics of the implemented biofeedback intervention. Different physiological systems were targeted, interventions were offered for varying durations of time, were applied both in a lab and real-life context, used different control conditions, and were operationalized as a stand-alone intervention or embedded in a broader intervention program. However, when plotting the occurrence of certain characteristics (such as duration) of a study against positive outcomes, visual inspection of the data did not reveal clear indications for better performance based on differences in this characteristic. Outcomes did not appear to improve when additional interventions were included, a real-life context was used or the duration was longer than 4 weeks.

Additionally, there were differences in the characteristics of the samples. The current review included both healthy subjects and individuals with health problems. There was no clear tendency for improved effectiveness of biofeedback in either group. It is difficult to compare the levels of stress between individuals and participant groups, since stress is a highly subjective experience and different assessment tools were used to measure stress. Dillon et al. (2016) induced stress in healthy participants prior to biofeedback and McCraty et al. (2009) reported that 21% of the sample showed high cortisol levels at baseline. However, the other

studies did not induce stress or report whether their samples displayed elevated stress at baseline. Individuals with severe stress-related problems might show more room for improvement on the outcome variables (e.g., HRV) as compared to participants with less severe problems. However, the current data do not allow assessing whether stress levels at baseline influenced the outcome of biofeedback.

Discussion

Elevated stress is a common negative psychological state that is associated with physiological changes. Therefore, stress management programs might benefit from the addition of a psychophysiological component. The purpose of this review was to investigate whether stress management interventions with a biofeedback component could improve both physiological and psychological indicators of stress. The results show that there is large diversity in the characteristics and effectiveness of the included interventions. No association between the occurrence of physiological and psychological effects could be observed. There is a need for further research assessing which components of biofeedback can improve stress outcomes.

Three studies observed significant improvements in both physiological and psychological stress indices in the biofeedback condition as compared to the control condition. Berry et al. (2014) implemented HRV coherence biofeedback in the context of chronic pain and achieved significantly higher HRV coherence and lower PSS scores after the intervention as compared to treatment as usual. The stress management study of Kotozaki et al. (2014) found that heart rate and cerebral blood flow biofeedback led to significantly decreased cortisol levels, decreased subjective stress, and neural changes. The mHealth design of Dillon et al. (2016) showed that EDA biofeedback games significantly decreased heart rate and perceived stress in comparison to a non-biofeedback game in a healthy population.

In addition to these three studies that observed effects on both outcome variables, six studies observed significantly stronger improvement in either physiological or psychological variables. HRV biofeedback significantly improves HRV in different participant populations, specifically individuals with stress-related chronic neck pain (Hallman et al. 2011), patients with coronary heart disease (Nolan et al. 2005), labourers (Sutarto et al. 2012), and students (Whited et al. 2014). Murphy (1984) showed that EMG biofeedback leads to significantly lower EMG values in a working class population. Nolan et al. (2005) and Sutarto et al. (2012) additionally observed reduced PSS and DASS scores after the intervention, but there was no significant difference with the control group. Lemaire et al. (2011) implemented stress management with HRV coherence biofeedback and

Table 4 Cross-tabulation of physiological and psychological outcomes of the biofeedback intervention as compared to the control condition

	Stress reduction	No reduction in stress	Total
Improvement in physiological parameters	3	5	8
No improvement in physiological parameters	1	5	6
Total	4	10	14

documented lower perceived stress levels that remained at 28 days follow-up.

The remaining five studies also observed improvement in some of the outcome variables, but group differences did not reach significance. Stress was targeted in white-collar workers by Allen and Blanchard (1980) and McCraty et al. (2009) through EMG and HRV coherence biofeedback. These studies recorded EMG and HRV reductions that did not differ from the control condition. No psychological effects were found. Mackay et al. (2015) and Siepman et al. (2014) on the other hand observed non-significant psychological effects, but no effects on breathing rate, muscle tension and heart rhythm, after implementing breathing rate and muscle tension biofeedback in multiple sclerosis patients or HRV biofeedback in pregnant women, respectively. Finally, in the study of Teufel et al. (2013) EDA biofeedback was associated with small to medium effects on EDA and small to large effects on perceived stress in an obese population, but no group differences could be found.

Several aspects of the study design could contribute to differences in the effectiveness of biofeedback. However, outcomes did not appear to vary according to specific intervention characteristics, such as duration, context or additional interventions. Sample characteristics could hamper statistical power to detect group differences. The studies included in the current review suffered from small sample sizes, sometimes reaching no more than ten participants per condition (Allen and Blanchard 1980; Berry et al. 2014; Teufel et al. 2013). Additionally, the majority of studies did not report whether the included sample showed elevated stress at baseline. Healthy individuals with few stress-related symptoms may not have a lot of room for improvement in the outcome measures. Finally, individual differences and preferences can also influence effectiveness through effects on motivation, engagement, and compliance (Marcus et al. 2012).

Biofeedback is based on the concept that increased control over physiological processes can improve performance and well-being. The current review assessed changes in both the physiological and psychological domain to better understand how improvements in these domains are related to one another. While Nolan et al. (2005) did find an inverse association between subjective stress and HRV modulation (the physiological outcome measure of the study) that was limited to the biofeedback condition, the current review did not observe a significant association between physiological and psychological outcomes variables. The mechanisms of change in biofeedback remain unclear. Berry et al. (2014) propose that biofeedback combined with self-regulation techniques can increase awareness of internal psychophysiological processes and hereby improve coping techniques (for pain). Biofeedback can not only increase awareness but also promote conscious control. Dillon et al. (2016) state that

stress can be caused by a perceived lack of control and that perceived control can be improved by teaching individuals how to master physiological processes through biofeedback.

The majority of included studies encouraged participants to practice the skills individually between sessions and two studies reported that over 90% of participants indeed performed (daily) exercises at home (Mackay et al. 2015; Whited et al. 2014). This finding is relevant for the development of mHealth applications using biofeedback since it suggests that participants are both able and motivated to apply biofeedback exercises individually and in different contexts. Benefits of an mHealth approach are that participants can use the intervention flexibly and at low cost, which can facilitate widespread adoption. A promising mHealth application for biofeedback is wearable technology, in which mobile sensory devices (e.g., wristbands) are used to non-invasively collect physiological data. Wearable devices can measure EDA (Villarejo et al. 2012), breathing patterns, and heart rhythm (and accordingly allow for the calculation of HRV) (Cropley et al. 2017). There is also a growing interest in using wearables to collect blood pressure and breathing rate as a guide for stress recovery training at home (through biofeedback), but further research is needed (e.g., Uddin et al. 2016). Ideally, however, a wide range of different physiological parameters could be collected by wearables, their data integrated and subsequently used as an additional source of information to guide preventive or therapeutic interventions (De Witte et al. 2018). There are some important practical barriers for implementation of mHealth, such as commercial availability of affordable and reliable physiological measurement tools, potential problems with data security, and diverse needs (for interfaces and tools) in different populations (Munos et al. 2016).

The current review was able to include one mHealth and game-based biofeedback intervention. Dillon et al. (2016) induced stress in their participants and subsequently offered EDA biofeedback through two mobile games. This approach has several potential advantages. It maximizes opportunities for generalization since participants learn to use biofeedback when they are stressed and the medium (a smartphone) makes it flexible to use in different contexts. Additionally, gamification elements could increase engagement and motivation. However, the effect sizes of Dillon et al. (2016) were modest and no follow-up measures were included. Other mobile biofeedback applications have been developed (e.g., Gaggioli et al. 2014; Munster-Segev et al. 2017), however, these applications still need to undergo rigorous testing and often lack gamification elements. Despite potential benefits, gamification is currently rarely included in stress management applications for adults (Hoffmann et al. 2017). Biofeedback video games have been developed for children and adolescents, such as Dojo and MindLight (Schoneveld et al. 2017).

Some limitations of the current review require discussion. Although the goal was to include all stress interventions using psychophysiological components, an exploratory literature search showed that this field of research was mostly limited to biofeedback interventions and the scope was consequently limited to increase comparability. A likely explanation for the lack of other interventions using psychophysiological measures is that, until recently, practical limitations (related to hardware or big data) impeded the possibilities for postponed feedback or report of psychophysiological data. As opposed to other interventions, biofeedback does not require large amounts of storage or computing power. Secondly, it is important to consider both the psychological and physiological outcome domains when investigating the effectiveness of psychophysiological interventions. Studies often did not investigate both outcome domains and could therefore not be included in the current review. The current review was only able to assess the association between these two outcome domains using dichotomous variables. Future research needs to investigate the association between physiological and psychological outcomes further by using analyses with more power, such as correlations. The included studies had very diverse samples. Comorbid conditions (such as obesity or chronic pain) could influence the effects of biofeedback. The large variability in the implemented control conditions and psychological measurement instruments did not allow to directly compare different studies and perform a meta-analysis. More high-quality standardized effectiveness studies of biofeedback interventions for stress are needed.

Taken together, the studies included in the current review suggest that biofeedback can be a promising intervention for stress management. It is feasible to explore the implementation of biofeedback in different contexts and in samples with stress associated to various conditions. One-third of the included studies, however, did not observe any incremental effectiveness of biofeedback. Small sample sizes and large differences between study designs also impede generalization. Additional research needs to determine which components of biofeedback are effective and establish whether psychophysiological measurements are an appropriate component of stress management programs. It is feasible to offer biofeedback as a gamified mHealth application and one included study supports that this is associated with positive short-term outcomes in both physiological and self-report indicators of stress.

Acknowledgements This work was written within the Carewear project, funded by a VLAIO TETRA grant (Grants IWT.150614 and HBC.2016.0099).

Compliance with Ethical Standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Allen, J. K., & Blanchard, E. B. (1980). Biofeedback-based stress management training with a population of business managers. *Biofeedback and Self-Regulation*, 5(4), 427–438.
- American Psychiatric Association. (2016). *Stress in America: the impact of discrimination*. Stress in America™ Survey.
- American Psychological Association. (2017). *Stress in America: The State of Our Nation*. Stress in America™ Survey.
- Bali, A., & Jaggi, A. S. (2015). Clinical experimental stress studies: methods and assessment. *Reviews in the Neurosciences*, 26(5), 555–579. <https://doi.org/10.1515/revneuro-2015-0004>.
- Berry, M. E., Chapple, I. T., Ginsberg, J. P., Gleichauf, K. J., Meyer, J. A., & Nagpal, M. L. (2014). Non-pharmacological Intervention for chronic pain in veterans: A pilot study of heart rate variability biofeedback. *Global Advances in Health and Medicine*, 3(2), 28–33. <https://doi.org/10.7453/gahmj.2013.075>.
- Chrousos, G. P., & Gold, P. W. (1992). The concepts of stress and stress system disorders. Overview of physical and behavioral homeostasis. *JAMA*, 267(9), 1244–1252.
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385–396.
- Cooper, C., & Dewe, P. (2008). Well-being—Absenteeism, presenteeism, costs and challenges. *Occupational Medicine*, 58(8), 522–524. <https://doi.org/10.1093/occmed/kqn124>.
- Cropley, M., Plans, D., Morelli, D., Sütterlin, S., Inceoglu, I., Thomas, G., & Chu, C. (2017). The association between work-related rumination and heart rate variability: A field study. *Frontiers in Human Neuroscience*, 11, 27. <https://doi.org/10.3389/fnhum.2017.00027>.
- De Witte, N. A. J., Bonroy, B., Debar, G., Sels, R., & Van Daele, T. (2018). *Carewear: An internet-based platform to implement wearable technology in mental health care*. Paper presented at the confere of the European Society for Research on Internet Interventions, Dublin.
- Dillon, A., Kelly, M., Robertson, I. H., & Robertson, D. A. (2016). Smartphone applications utilizing biofeedback can aid stress reduction. *Frontiers in Psychology*, 7, 832. <https://doi.org/10.3389/fpsyg.2016.00832>.
- Gaggioli, A., Cipresso, P., Serino, S., Campanaro, D. M., Pallavicini, F., Wiederhold, B. K., & Riva, G. (2014). Positive technology: A free mobile platform for the self-management of psychological stress. *Studies in Health Technology and Informatics*, 199, 25–29.
- Hallman, D. M., Olsson, E. M., von Schéele, B., Melin, L., & Lyskov, E. (2011). Effects of heart rate variability biofeedback in subjects with stress-related chronic neck pain: A pilot study. *Applied Psychophysiology and Biofeedback*, 36(2), 71–80. <https://doi.org/10.1007/s10484-011-9147-0>.
- Higgins, J. P., Altman, D. G., Gøtzsche, P. C., Jüni, P., Moher, D., Oxman, A. D., ... Group, C. S. M. (2011). The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*, 343, d5928. <https://doi.org/10.1136/bmj.d5928>.
- Hoffmann, A., Christmann, C. A., & Bleser, G. (2017). Gamification in stress management apps: A critical app review. *JMIR Serious Games*, 5(2), e13. <https://doi.org/10.2196/games.7216>.
- Istepanian, R., Jovanov, E., & Zhang, Y. T. (2004). Introduction to the special section on M-Health: Beyond seamless mobility and

- global wireless health-care connectivity. *IEEE Transactions on Information Technology in Biomedicine*, 8(4), 405–414.
- Jarczok, M., Jarczok, M., Mauss, D., Koenig, J., Li, J., Herr, R., & Thayer, J. (2013). Autonomic nervous system activity and workplace stressors—A systematic review. *Neuroscience and Biobehavioral Reviews*, 37(8), 1810–1823. <https://doi.org/10.1016/j.neubiorev.2013.07.004>.
- Kotozaki, Y., Takeuchi, H., Sekiguchi, A., Yamamoto, Y., Shinada, T., Araki, T., ... Kawashima, R. (2014). Biofeedback-based training for stress management in daily hassles: An intervention study. *Brain and Behavior*, 4(4), 566–579. <https://doi.org/10.1002/brb3.241>.
- Lemaire, J. B., Wallace, J. E., Lewin, A. M., de Groot, J., & Schaefer, J. P. (2011). The effect of a biofeedback-based stress management tool on physician stress: A randomized controlled clinical trial. *Open Medicine*, 5(4), e154–e163.
- Lin, H., Lin, H., Lin, W., & Huang, A. (2011). Effects of stress, depression, and their interaction on heart rate, skin conductance, finger temperature, and respiratory rate: Sympathetic-parasympathetic hypothesis of stress and depression. *Journal of Clinical Psychology*, 67(10), 1080–1091. <https://doi.org/10.1002/jclp.20833>.
- Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the depression anxiety stress scales*, 2nd ed. Sydney: Psychology Foundation.
- Mackay, A. M., Buckingham, R., Schwartz, R. S., Hodgkinson, S., Beran, R. G., & Cordato, D. J. (2015). The effect of biofeedback as a psychological intervention in multiple sclerosis: A randomized controlled study. *International Journal of MS Care*, 17(3), 101–108. <https://doi.org/10.7224/1537-2073.2014-006>.
- Marcus, S. M., Stuart, E. A., Wang, P., Shadish, W. R., & Steiner, P. M. (2012). Estimating the causal effect of randomization versus treatment preference in a doubly randomized preference trial. *Psychological Methods*, 17(2), 244–254. <https://doi.org/10.1037/a0028031>.
- McCraty, R., Atkinson, M., Lipsenthal, L., & Arguelles, L. (2009). New hope for correctional officers: An innovative program for reducing stress and health risks. *Applied Psychophysiology and Biofeedback*, 34(4), 251–272. <https://doi.org/10.1007/s10484-009-9087-0>.
- Munos, B., Baker, P. C., Bot, B. M., Crouthamel, M., de Vries, G., Ferguson, I., ... Wang, P. (2016). Mobile health: The power of wearables, sensors, and apps to transform clinical trials. *Annals of the New York Academy of Sciences*, 1375(1), 3–18. <https://doi.org/10.1111/nyas.13117>.
- Munster-Segev, M., Fuerst, O., Kaplan, S. A., & Cahn, A. (2017). Incorporation of a stress reducing mobile app in the care of patients with type 2 diabetes: A prospective study. *JMIR mHealth uHealth*, 5(5), e75. <https://doi.org/10.2196/mhealth.7408>.
- Murphy, L. R. (1984). Stress management in highway maintenance workers. *Journal of Occupational Medicine*, 26(6), 436–442.
- Nolan, R. P., Kamath, M. V., Floras, J. S., Stanley, J., Pang, C., Picton, P., & Young, Q. R. (2005). Heart rate variability biofeedback as a behavioral neurocardiac intervention to enhance vagal heart rate control. *American Heart Journal*, 149(6), 1137. <https://doi.org/10.1016/j.ahj.2005.03.015>.
- Schoenberg, P. L., & David, A. S. (2014). Biofeedback for psychiatric disorders: A systematic review. *Applied Psychophysiology and Biofeedback*, 39(2), 109–135. <https://doi.org/10.1007/s10484-014-9246-9>.
- Schoneveld, E. A., Lichtwarck-Aschoff, A., & Granic, I. (2017). Preventing childhood anxiety disorders: Is an applied game as effective as a cognitive behavioral therapy-based program? *Prevention Science*. <https://doi.org/10.1007/s11121-017-0843-8>.
- Siepmann, M., Hennig, U. D., Siepmann, T., Nitzsche, K., Mück-Weymann, M., Petrowski, K., & Weidner, K. (2014). The effects of heart rate variability biofeedback in patients with preterm labour. *Applied Psychophysiology and Biofeedback*, 39(1), 27–35. <https://doi.org/10.1007/s10484-013-9238-1>.
- Sutarto, A. P., Wahab, M. N., & Zin, N. M. (2012). Resonant breathing biofeedback training for stress reduction among manufacturing operators. *International Journal of Occupational Safety and Ergonomics*, 18(4), 549–561. <https://doi.org/10.1080/10803548.2012.11076959>.
- Teufel, M., Stephan, K., Kowalski, A., Käsberger, S., Enck, P., Zipfel, S., & Giel, K. E. (2013). Impact of biofeedback on self-efficacy and stress reduction in obesity: A randomized controlled pilot study. *Applied Psychophysiology and Biofeedback*, 38(3), 177–184. <https://doi.org/10.1007/s10484-013-9223-8>.
- Uddin, A. A., Morita, P. P., Tallevi, K., Armour, K., Li, J., Nolan, R. P., & Cafazzo, J. A. (2016). Development of a wearable cardiac monitoring system for behavioral neurocardiac training: A usability study. *JMIR mHealth uHealth*, 4(2), e45. <https://doi.org/10.2196/mhealth.5288>.
- Van Daele, T., Hermans, D., Van Audenhove, C., & Van den Bergh, O. (2012). Stress reduction through psychoeducation: A meta-analytic review. *Health Education & Behavior*, 39(4), 474–485. <https://doi.org/10.1177/1090198111419202>.
- Villarejo, M. V., Zapirain, B. G., & Zorrilla, A. M. (2012). A stress sensor based on Galvanic Skin Response (GSR) controlled by ZigBee. *Sensors (Basel)*, 12(5), 6075–6101. <https://doi.org/10.3390/s120506075>.
- Wheat, A. L., & Larkin, K. T. (2010). Biofeedback of heart rate variability and related physiology: A critical review. *Applied Psychophysiology and Biofeedback*, 35(3), 229–242. <https://doi.org/10.1007/s10484-010-9133-y>.
- Whited, A., Larkin, K. T., & Whited, M. (2014). Effectiveness of emWave biofeedback in improving heart rate variability reactivity to and recovery from stress. *Applied Psychophysiology and Biofeedback*, 39(2), 75–88. <https://doi.org/10.1007/s10484-014-9243-z>.
- Williams, A. D. (2016). *Harnessing the quantified self-movement for optimal mental health and wellbeing*. Paper presented at the LTA, Amsterdam.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.